

Results from $\Upsilon(5S)$ at Belle: Strange Beauty and other Beasts*

K. Kinoshita

University of Cincinnati
PO Box 210011
Cincinnati, OH 45221, USA
E-mail: kay.kinoshita@uc.edu

Abstract

The Belle experiment collected in 2005-6 a total of 23.6 fb^{-1} of data at the $\Upsilon(10860)$ resonance, also known as $\Upsilon(5S)$. These constitute nearly all of the world's sample of $e^+e^- \rightarrow \Upsilon(10860)$ events, which provide clean B_s pairs. We present here several B_s and $\Upsilon(10860)$ properties recently extracted from these data.

* talk presented at the Lake Louise Winter Institute, 18-23 February, 2008

Belle,[1] one of two B -factory experiments, is located at KEKB[2] and studies primarily the CP asymmetries of B meson decay in e^+e^- annihilations at the $\Upsilon(4S)$ resonance ($Mc^2 = 10580 \pm 1$ MeV, $\Gamma = 20.5 \pm 2.5$ MeV).[3] However, the richness of Belle physics reaches well beyond CP (see, e.g., <http://belle.kek.jp/>); an unequalled sample of $B\bar{B}$, $c\bar{c}$, $\tau\bar{\tau}$, and two-photon events enables access to many other topics. We now add B_s and bottomonium studies, thanks to data collected at the $\Upsilon(10860)$ resonance.

The $\Upsilon(10860)$, ($Mc^2 = 10865 \pm 8$ MeV/ c^2 , $\Gamma = 110 \pm 13$ MeV),[3] is interpreted as $\Upsilon(5S)$, the fourth excitation of the vector bound state of $b\bar{b}$. It is above $B_s\bar{B}_s$ threshold and, given the success of the $\Upsilon(4S)$ program in characterizing properties of $B_{d,u}$, it is natural to contemplate B_s at $\Upsilon(10860)$. The e^+e^- environment produces clean events, efficiently triggered, with precisely known center-of-mass energy. Furthermore, the B -factory offers an existing facility with high luminosity, a well-studied detector, and $\Upsilon(4S)$ data for comparisons. On the other hand, high integrated luminosities are needed, $\sigma(e^+e^- \rightarrow \Upsilon(10860)) \approx \sigma(e^+e^- \rightarrow \Upsilon(4S))/3 \approx 0.3$ nb, and events include $B_{d,u}$ as well as B_s . While B_s is produced copiously in hadronic collisions, the $\Upsilon(10860)$ can be competitive, particularly in aspects of B_s decay that are limited by systematic effects at a hadron machine.

Belle has run twice at the $\Upsilon(10860)$. In June 2005 a three-day “engineering” run served to test KEKB, which had never operated at energies above the $\Upsilon(4S)$, and study the basics of $\Upsilon(10860)$, B_s , and B_s^* . A scan of five energy points was used to locate the peak, at $\sqrt{s} = 10869$ MeV, where 1.86 fb^{-1} were collected. By June 2006, results confirmed the projected potential of $\Upsilon(10860)$, and 21.7 fb^{-1} were collected in 20 days.

As prerequisite to studies of B_s , its abundance in $\Upsilon(10860)$ events was determined from the 2005 data.[4] About 10% of the hadronic events are resonance (assumed to be $b\bar{b}$), the rest being continuum $e^+e^- \rightarrow q\bar{q}$ ($q = u, d, s, c$). Figure 1 (L) displays R_2 , the ratio of the 2nd and 0th Fox-Wolfram moments,[5] a measure of “jettiness” that tends to be lower for the more isotropic resonance events. We find $(3.01 \pm 0.02 \pm 0.16) \times 10^5 \text{ } b\bar{b} \text{ events/fb}^{-1}$. The $b\bar{b}$ events may fragment to: $B_s^{(*)}\bar{B}_s^{(*)}$, $B_q^{(*)}\bar{B}_q^{(*)}$, $B_q\bar{B}_q^{(*)}\pi$, $B_q\bar{B}_q\pi\pi$ (q is a u - or d -quark). To determine the fraction (f_s) that are $B_s^{(*)}\bar{B}_s^{(*)}$, we measure the inclusive rate $\mathcal{B}(\Upsilon(10860) \rightarrow D_s X) \equiv \mathcal{B}_\Upsilon$, an average over B_s , B_d , and B_u (Figure 1(R)). Measured $\mathcal{B}(B_{u,d} \rightarrow D_s X)$ and our understanding of decay mechanisms are used to estimate $\mathcal{B}(B_s \rightarrow D_s X) = (92 \pm 11)\%$. [6] Assuming that B_d and B_u are produced equally, $\mathcal{B}_\Upsilon$ depends only on f_s . The same analysis, performed with inclusive D^0 yields, gives an independent value of f_s with larger uncertainties; $\mathcal{B}(B_s \rightarrow D^0 X) \ll \mathcal{B}(B_q \rightarrow D^0 X)$. The combined result is $f_s = (18.0 \pm 1.3 \pm 3.2)\%$. [4]

The results below, based on 23.6 fb^{-1} , cover $B_s \rightarrow D_s\pi$, $D_s K$ (preliminary), $\Upsilon(10860) \rightarrow \Upsilon(nS)\pi^+\pi^-$, [8] and $B_s \rightarrow \gamma\gamma$, $\phi\gamma$. [9] B_s decays are identified by “full reconstruction” of all final state particles, a method used with great success at the $\Upsilon(4S)$. Each candidate’s energy and momentum in the e^+e^- center-of-mass are evaluated as $\Delta E \equiv E_{cand} - E_{beam}$ and $M_{bc} \equiv \sqrt{E_{beam}^2 - p_{cand}^2}$. In $B_s\bar{B}_s$ (analogous to $\Upsilon(4S) \rightarrow B_q\bar{B}_q$), the B_s carries the beam energy, so $\langle \Delta E \rangle = 0$ and $\langle M_{bc} \rangle = M_{B_s}$. For $B_s^*\bar{B}_s$ or $B_s^*\bar{B}_s^*$, the kinematics of $B_s^* \rightarrow B_s\gamma$ (with 50 MeV energy) leads to localized signals as well. In the case of $B_s^*\bar{B}_s$ the result is effectively the loss of 50 MeV from the B_s pair, with the energy being shared approximately equally: $\langle \Delta E \rangle \approx -25$ MeV and $\langle M_{bc} \rangle \approx M_{B_s} + 25$ MeV. For $B_s^*\bar{B}_s^*$ the energy reduction is ~ 50 MeV.

The decay $B_s \rightarrow D_s^-\pi^+$ proceeds dominantly via a CKM-favored spectator process. D_s is reconstructed in $\phi(\rightarrow K^+K^-)\pi^-$, $K^{*0}(\rightarrow K^+K^-)K^-$, and $K_S(\rightarrow \pi^+\pi^-)K^-$. Shown in Figure 2(L) is the distribution in ΔE and M_{bc} for data, with signal boxes for $B_s^*\bar{B}_s^*$,




FIG. 1: (Left) Distribution in R_2 , (histogram) 1.86 fb^{-1} at the $\Upsilon(10860)$ and (points) continuum below the $\Upsilon(4S)$, scaled. (Right) Yield of D_s mesons as a function of $x \equiv p_{D_s} / \sqrt{E_{beam}^2 - M_{D_s}^2}$, (points) $\Upsilon(10860)$ and (histogram) scaled continuum.




FIG. 2: Distributions in ΔE and M_{bc} of B_s candidates in 23.6 fb^{-1} of Belle data at $\Upsilon(10860)$. (Left) $B_s \rightarrow D_s \pi$, with signal regions for $B_s^* \bar{B}_s^*$ (upper signal box), $B_s^* \bar{B}_s$ (middle box), and $B_s \bar{B}_s$ (lower box) events. (Right) $B_s \rightarrow D_s K$, with signal region for $B_s^* \bar{B}_s^*$ (solid box) and region accumulating backgrounds from $D_s \pi$ (dashed box).

$B_s^* \bar{B}_s$ (inclusion of $B_s \bar{B}_s^*$ is implied), and $B_s \bar{B}_s$ events. A two-dimensional fit yields $B_s \rightarrow D_s \pi$ signals of 147_{-13}^{+14} (20σ), $12.7_{-5.1}^{+6.0}$ (2.9σ), and $3.8_{-3.6}^{+4.5}$ (1.1σ) in the $B_s^* \bar{B}_s^*$, $B_s^* \bar{B}_s$, and $B_s \bar{B}_s$ channels, respectively. Using $f_s = (19.5_{-2.3}^{+3.0})\%$ and $\sigma_{e^+e^- \rightarrow b\bar{b}} = 0.302 \pm 0.014 \text{ nb}$ (a weighted average from [4, 7]), we obtain $\mathcal{B} = (3.41_{-0.31}^{+0.33}(\text{stat.})_{-0.41}^{+0.42}(\text{sys.})_{-0.45}^{+0.46}(f_s)_{-0.28}^{+0.33}(\mathcal{B}(D_s^- \rightarrow \phi\pi))) \times 10^{-3}$ and $f_{B_s^* B_s^*} \equiv \frac{\sigma(e^+e^- \rightarrow B_s^* \bar{B}_s^*)}{\sigma(e^+e^- \rightarrow B_s^* \bar{B}_s^*) + \sigma(e^+e^- \rightarrow B_s^* \bar{B}_s)} = (89.8_{-4.0}^{+3.8})\%$, $f_{B_s^* B_s} \equiv \frac{\sigma(e^+e^- \rightarrow B_s^* \bar{B}_s + B_s \bar{B}_s^*)}{\sigma(e^+e^- \rightarrow B_s^* \bar{B}_s^*)} = (7.8_{-3.0}^{+3.3})\%$. The masses are found from the peak locations, $m_{B_s} = (5364.8 \pm 1.3 \pm 2.4) \text{ MeV}/c^2$ and $m_{B_s^*} = (5417.6 \pm 0.4 \pm 0.5) \text{ MeV}/c^2$.

The same process, CKM-suppressed, leads to $B_s \rightarrow D_s^- K^+$. Given the lower expected rate, only $B_s^* \bar{B}_s^*$ is examined. Kinematic similarities result in a large incursion of $B_s \rightarrow D_s^- \pi^+$ into the signal region (Figure 2(R)). Good hadron identification is thus crucial. A two-dimensional fit yields $6.8_{-2.7}^{+3.4}$ (3.6σ) events. We obtain $\mathcal{B} = (2.2_{-0.9}^{+1.1}(\text{stat.}) \pm 0.3(\text{sys.}) \pm 0.3(f_s) \pm 0.2(\mathcal{B}(D_s^- \rightarrow \phi\pi))) \times 10^{-4}$ and $\frac{\mathcal{B}(B_s \rightarrow D_s K)}{\mathcal{B}(B_s \rightarrow D_s \pi)} = (6.6_{-2.8}^{+3.4})\%$.

The observation of $Y(4260)$, [10] and similar charmonium-like particles, [11] raises the question of whether analogous bottomonium-like particles exist in the region near and above $\Upsilon(10860)$. [12] The Y 's are observed in modes $\psi(nS)h^+h^-$ where h is π or K ; their nature is not yet fully understood. Given widths of order 10 keV for $\Upsilon(nS) \rightarrow \Upsilon(1S)\pi^+\pi^-$

($n = 2, 3, 4$), one would naïvely expect $\mathcal{B}(\Upsilon(5S) \rightarrow \Upsilon(1S)\pi^+\pi^-)$ to be tiny, of order 10^{-5} . Belle has searched for $\Upsilon(10860) \rightarrow \Upsilon(nS)(\rightarrow \mu^+\mu^-)h^+h^-$. Figure 3(L) shows the distribution of $\mu^+\mu^-$ mass and $\Delta M \equiv M(\mu\mu\pi\pi) - M(\mu\mu)$. The horizontal bands indicate signal regions for $\Upsilon(1S/2S/3S) \rightarrow \mu\mu$. Peaks corresponding to $\Upsilon(10860) \rightarrow \Upsilon(nS)\pi^+\pi^-$ are seen in projections onto ΔM of Υ signal bands (Figure 3(R)) which are fitted to determine signal yields. From these we find cross sections, branching fractions, and partial widths for four modes, assuming $\Upsilon(10860) = \Upsilon(5S)$: (in MeV) $\Gamma(\Upsilon(5S) \rightarrow \Upsilon(1S)\pi^+\pi^-) = 0.59 \pm 0.04 \pm 0.09$, $\Gamma(\Upsilon(5S) \rightarrow \Upsilon(2S)\pi^+\pi^-) = 0.85 \pm 0.07 \pm 0.16$, $\Gamma(\Upsilon(5S) \rightarrow \Upsilon(3S)\pi^+\pi^-) = 0.52^{+0.20}_{-0.17} \pm 0.10$, $\Gamma(\Upsilon(5S) \rightarrow \Upsilon(1S)K^+K^-) = 0.067^{+0.017}_{-0.015} \pm 0.013$. The $\Upsilon\pi\pi$ widths are of order 1 MeV.




FIG. 3: (Left) Distribution in data of $M(\mu^+\mu^-)$ vs. $\Delta M \equiv M(\mu\mu\pi\pi) - M(\mu\mu)$ of $\mu^+\mu^-\pi^+\pi^-$ combinations. The line indicates the kinematic limit, $M(\mu\mu\pi\pi) = \sqrt{s}$. The horizontal bands indicate signal regions for (bottom to top) $\Upsilon(1S)$, $\Upsilon(2S)$, and $\Upsilon(3S)$. (Right) Projections onto ΔM of (top) $\Upsilon(2S)$ and (bottom) $\Upsilon(1S)$ signal bands.

The widths for $\Upsilon(2S/3S/4S/10860) \rightarrow \Upsilon(1S)\pi^+\pi^-$ [3, 8, 13] are 0.0060, 0.0009, 0.0019, and 0.59 MeV. The rate for $\Upsilon(10860)$ is largest by two orders of magnitude. While the rate for $\Upsilon(5S)$ could be substantially enhanced,[14] our result raises the possibility that the $\Upsilon(10860)$ is not a pure $\Upsilon(5S)$ state. To address this issue, Belle has scanned in energy in the region above the $\Upsilon(10860)$ resonance and anticipates first results by summer 2008.

Belle has searched for $B_s \rightarrow \gamma\gamma$ and $B_s \rightarrow \phi\gamma$. [9] Each includes a hard photon, readily isolated in the B -factory but difficult to trigger on at a hadron machine. For the Standard Model, $\mathcal{B}(B_s \rightarrow \gamma\gamma)$ is in the range $(0.4 - 1.0) \times 10^{-6}$; extended models give rates up to 5×10^{-6} . The Belle $\Upsilon(10860)$ data improve the limit 20-fold over previous searches, $\mathcal{B} < 8.7 \times 10^{-6}$ (90% CL). The process for $B_s \rightarrow \phi\gamma$ is identical to that for $B \rightarrow K^{(*)}\gamma$ ($\mathcal{B} \sim 4 \times 10^{-5}$) and is expected at this level. We report the first observation, $\mathcal{B}(B_s \rightarrow \phi\gamma) = (57^{+18}_{-15}(stat)^{+12}_{-11}(sys)) \times 10^{-6}$, with a significance of 5.5σ .

In summary, Belle, which was designed for precision measurements of CKM parameters, has opened a new avenue of inquiry by collecting 23.6 fb^{-1} at the $\Upsilon(10860)$ resonance. The sample includes ≈ 1.3 million B_s events and has yielded unprecedented sensitivity and precision for several modes. In addition, an anomaly in the rate of $\Upsilon(10860) \rightarrow \Upsilon(nS)h^+h^-$ is stimulating new interest in bottomonium studies.

Acknowledgments

The author wishes to thank the LLWI organizers and staff. This work is supported by Department of Energy grant # DE-FG02-84ER40153.

-
- [1] Belle Collaboration, A. Abashian *et al.*, Nucl. Instr. Meth. A **479**, 117 (2002); Z.Natkaniec *et al.* (Belle SVD2 Group), Nucl. Instr. Meth. A **560**, 1(2006).
 - [2] S. Kurokawa and E. Kikutani, Nucl. Instrum. Meth., **A499**, 1 (2003), and other papers included in this Volume.
 - [3] W.-M. Yao *et al.*, J. Phys., **G 33**, 1 (2006), 2007 partial update for 2008
 - [4] A. Drutskoy *et al.* (Belle Collaboration), Phys. Rev. Lett. **98**, 052001 (2007).
 - [5] G.C. Fox and S. Wolfram, Phys. Rev. Lett. **41**, 1581 (1978).
 - [6] M. Artuso *et al.* (CLEO Collaboration), Phys. Rev. Lett. **95**, 261801 (2005).
 - [7] G.S. Huang *et al.* (CLEO Collaboration), Phys. Rev. **D 75**, 012002 (2007).
 - [8] K.-F. Chen, W.-S. Hou, M. Shapkin, A. Sokolov, *et al.* (Belle Collaboration), Phys. Rev. Lett. **100**, 112001 (2008).
 - [9] J. Wicht *et al.* (Belle Collaboration), Phys. Rev. Lett. **100**, 121801 (2008).
 - [10] B. Aubert *et al.* (Babar Collaboration), Phys. Rev. Lett. **95**, 142001 (2005); Q. He *et al.* (CLEO Collaboration), Phys. Rev. **D 74**, 091104 (R) (2006); C.Z. Yuan *et al.* (Belle Collaboration), Phys. Rev. Lett. **99**, 182004 (2007).
 - [11] B. Aubert *et al.* (Babar), Phys. Rev. Lett. **98**, 212001 (2007); X.L. Wang *et al.* (Belle), **99**, 142002 (2007); T.E. Coan *et al.* (CLEO), **96**, 162003 (2006); C.Z. Yuan, C.P. Shen, P. Wang, X.L. Wang *et al.* (Belle), Phys. Rev. **D 77**, 011105 (R) (2008).
 - [12] W.S. Hou, Phys. Rev. **D 74**, 017504 (2006).
 - [13] B. Aubert *et al.* (Babar Collaboration), Phys. Rev. Lett. **96**, 232001 (2006); A. Sokolov *et al.* (Belle Collaboration), Phys. Rev. **D 75**, 071103 (2007).
 - [14] Yu. A. Siminov, JETP Lett. **87**, 147 (2008); C. Meng and K.-T. Chao, Phys. Rev. **D 77**, 074003 (2008).